

DOI: 10.22616/foodbalt.2017.019









- variation in phenolic composition and antioxidant and anti-inflammatory activities of *Calamintha nepeta* (L.) Savi. *Food Research International*, Vol. 69, p. 121–132.
25. Prasad K.N., Yang B., Yang S., Chen Y., Zhao M., Ashraf M., Jiang Y. (2009) Identification of phenolic compounds and appraisal of antioxidant and antityrosinase activities from litchi (*Litchi sinensis* Sonn.) seeds. *Food Chemistry*, Vol. 116(1), p. 1–7.
26. Raghavan Uhl S. (2000) *Handbook of Spices, Seasonings, and Flavorings*. Boca Raton: Technomic Publishing, Lancaster, 352 p.
27. Re R., Pellegrini N., Proteggente A., Pannala A., Yang M., Rice-Evans C. (1999) Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Radical Biology and Medicine*, Vol. 26, p. 1231–1237.
28. Singh N., Rajini P. S. (2004) Free radical scavenging activity of an aqueous extract of potato peel. *Food Chemistry*, Vol. 85, p. 611–616.
29. Singleton V.L., Orthofer R., Lamuela-Raventos R.M. (1999) Analysis of total phenols and other oxidation substrates and antioxidants by means of Folin-Ciocalteu reagent. *Methods in Enzymology*, Vol. 29, p. 152–178.
30. Tomsone L., Kruma Z., Lepse L. (2012) Influence of genotype and harvest time on the phenolic content of horseradish (*Armoracia rusticana* L.) roots. In: Research for rural development 2012: Annual 18th international scientific conference proceedings, Jelgava, 16-18 May 2012 / Latvia University of Agriculture. Jelgava: LLU, Vol. 1, p. 124–130.
31. Zhou K., Yu L. (2006) Total phenolic contents and antioxidant properties of commonly consumed vegetables grown in Colorado, *LWT - Food Science and Technology*, Vol. 39, p. 1155–1162.
32. Xing R., Yu H. H., Liu S., Zhang W. W., Zhang Q. B., Li Z. E. (2005) Antioxidant activity of differently regioselective chitosan sulfates in vitro. *Bioorganic & Medicinal Chemistry*, Vol. 13, p. 1387–1392.
33. Yu L., Haley S., Perret J., Harris M., Wilson J., Haley S. (2003) Antioxidant properties of bran extracts from Akron wheat grown at different locations. *Journal of Agriculture and Food Chemistry*, Vol. 51, p. 1566–1570.